



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Rusts of Oregon.—JACKSON²⁰ has published an annotated list of the rusts of Oregon, which brings together for the first time the rust flora of a state on the Pacific coast. All of the grain rusts recorded for North America (except *Puccinia Sorghi*) are known to occur in the state, and also all of the rusts of greenhouse crops. In addition to these, the Pacific coast rust of pears and quinces is said to be of considerable economic importance; and of course the forest-tree rusts represent an important field of investigation. The list includes 220 species of rusts occurring on about 500 different hosts, 8 of the species being described as new.—J. M. C.

Practical breeding.—COLLINS and KEMPTON²¹ have given an excellent example of the effective application of the principles of pure science to the solution of a practical problem. The production of a race of sweet corn resistant to the earworm has been a strictly practical problem, and introduces no new phenomena or theories of inheritance. The authors, however, have established statistically the correlation between the amount of damage done by the earworm and certain superficial plant characters, and have followed this by selective breeding for those significant characters.—MERLE C. COULTER.

The morning glory in genetics.—BARKER²² has found that the morning-glory is very favorable material for work in genetics. The almost innumerable combinations of floral colors are beautifully explained by the enzyme theory. "Each epistatic type is due to the addition of one or more genes, probably enzymatic in nature, which are not present in the hypostatic type."—MERLE C. COULTER.

Rusts of Cuba.—ARTHUR and JOHNSTON²³ have brought together all collections of Cuban rusts as a "basis for a thoroughly scientific and economic exploration of the island." The list includes 140 species, 12 of which are described as new, 15 are new to the North American flora, and 10 are exclusively Cuban.—J. M. C.

²⁰ JACKSON, H. S., The Uredinales of Oregon. Mem. Brooklyn Bot. Gard. 1:198-297. 1918.

²¹ COLLINS, G. N., and KEMPTON, J. H., Breeding sweet corn resistant to the corn earworm. Jour. Agric. Research 12:549-572. 1917.

²² BARKER, E. E., Hereditary studies in the morning-glory (*Ipomaea purpurea*). Cornell Univ. Agric. Exper. Sta. Bull. no. 392. pp. 38. pls. 3. 1917.

²³ ARTHUR, J. C., and JOHNSTON, J. R., Uredinales of Cuba. Mem. Torr. Bot. Club 17:97-175. 1918.